arXiv:1512.03552v1 [math.PR] 11 Dec 2015

REGULARITY OF THE DRIFT AND ENTROPY OF RANDOM WALKS ON GROUPS

LORENZ GILCH AND FRANÇOIS LEDRAPPIER

Random walks on a group G model many natural phenomena. A random walk is defined by a probability measure p on G. We are interested in asymptotic properties of the random walks and in particular in the linear drift and the asymptotic entropy. If the geometry of the group is rich, then these numbers are both positive and the way of dependence on p is itself a property of G. In this note, we review recent results about the regularity of the drift and the entropy for free groups, free products and hyperbolic groups.

1. Entropy and linear drift

We recall in this section the main notations for the objects under consideration associated to a group G and a probability measure p on G. Background on random walks can be found in the survey papers [**KV**] and [**V**] and in the book by W. Woess ([**W**]).

Let G be a finitely generated group and S a symmetric finite generating set. For $g \in G$, let $|g|$ denote the smallest $n \in \mathbb{N}$ such that g can be written as $g = s_1 \cdots s_n$, where $s_1, \ldots, s_n \in S$. We denote by $d(g,h) := |g^{-1}h|$ the left invariant associated metric. Let p be a probability measure on G with support B. Unless otherwise specified, we always assume that B is finite and that $\bigcup_{n\in\mathbb{N}} B^n = G$. We denote by $\mathcal{P}(B)$ the set of probability measures with support B. The set $\mathcal{P}(B)$ is naturally identified with an open subset of the probabilities on B, which is a contractible open polygonal bounded convex domain in $\mathbb{R}^{|B|-1}$. We form, with $p^{(0)}$ being the Dirac measure at the identity e,

$$p^{(n)}(g) \;=\; [p^{(n-1)} \star p](g) \;=\; \sum_{h\in G} p^{(n-1)}(gh^{-1})p(h),$$

where $g \in G$. Let $\varrho(p)$ be the spectral radius of the Markov operator on $\ell_2(G)$ defined by p. It is given by $\varrho(p) = \limsup_{n\to\infty} p^{(n)}(e)^{1/n}$. Define the entropy $H_{n,p}$ and the drift $L_{n,p}$ of $p^{(n)}$ by:

$$H_{n,p} := -\sum_{g\in G} p^{(n)}(g)\ln p^{(n)}(g), \quad L_{n,p} := \sum_{g\in G} |g| p^{(n)}(g),$$

and the asymptotic entropy h_p and the linear drift ℓ_p by

$$h_p := \lim_{n\to\infty} \frac{1}{n} H_{n,p}, \quad \ell_p := \lim_{n\to\infty} \frac{1}{n} L_{n,p}.$$

2000 *Mathematics Subject Classification.* Primary 60J10; Secondary 28D20, 20E06.
Key words and phrases. random walks, drift, entropy, analytic.

Both limits exist by subadditivity and Fekete's Lemma. The linear drift makes sense as soon as $\sum_{g\in G}|g|p(g)<+\infty$, the entropy under the slightly weaker condition $H_{1,p}<+\infty$. The entropy h_p was introduced by Avez ([**Av**]) and is related to bounded solutions of the equation on G of the form $f(g)=\sum_{h\in G}f(gh)p(h)$ (see e.g. [**KV**]). In particular, $h_p=0$ if and only if the only bounded solutions (called *harmonic* functions) are the constant functions ([**KV**], [**De2**]). A general relation is ([**Gu**])

$$h_p \;\leq\; \ell_p v, \tag{1}$$

where $v:=\lim_{n\to\infty}\frac{1}{n}\ln\left(\#\{g\in G;|g|\leq n\}\right)$ is the *volume entropy* of G. In particular, if $\ell_p=0$ then $h_p=0$.

We say that p is symmetric if $B=B^{-1}$ and $p(g)=p(g^{-1})$ for all $g\in B$. We call p centered if $\sum_{g\in B}\chi(g)p(g)=0$ for all group morphisms $\chi:G\to\mathbb{R}$. Clearly, symmetric probabilities are centered. If p is centered and $h_p=0$, then $\ell_p=0$ ([**Va**], [**Ma1**]). If p is not centered, we may have $h_p=0$ and $\ell_p\neq 0$, for instance on $\mathbb{Z}$. If this is the case, there is a group morphism $\chi:G\to\mathbb{R}$ such that $\ell_p=\sum_{g\in B}\chi(g)p(g)$ ([**KL**], see also [**EKn**] for finite versions of this result).

Both h_p and ℓ_p describe asymptotic properties of the random walk directed by p. Let $(\Omega,P)=(G^{\mathbb{N}},p^{\otimes\mathbb{N}})$ be the infinite product space so that $\omega=(\omega_1,\omega_2,\ldots)\in G^{\mathbb{N}}$ is realized by a sequence of i.i.d. random variables with values in G and distribution p. We form the right random walk by $X_n(\omega):=\omega_1\omega_2\cdots\omega_n$. The probability $p^{(n)}$ is the distribution of X_n, and an application of Kingman's subadditive ergodic theorem ([**Ki**]) gives that, for P-a.e. ω,

$$\lim_{n\to\infty}\frac{1}{n}|X_n| \;=\; \ell_p \quad\text{and}\quad \lim_{n\to\infty}-\frac{1}{n}\ln\left(p^{(n)}(X_n)\right) \;=\; h_p. \tag{2}$$

The random walk is said to be recurrent if, for P-a.e. ω, there is a positive $n\in\mathbb{N}$ with $X_n(\omega)=e$. In this case there is an infinite number of integers n with $X_n=e$ and, by (2), $\ell_p=0$. Hence, $h_p=0$. From here on, we assume that the random walk is transient, i.e. $|X_n|\to\infty$ for P-a.e. ω. The *Green function* $G(g,h)$, $g,h\in G$, is defined by

$$G(g,h) \;:=\; \sum_{n\geq 0}p^{(n)}(g^{-1}h).$$

By decomposing of the first visit to h and using transitivity of the random walk we get

$$G(g,h) \;=\; F(g,h)G(h,h) \;=\; F(g,h)G(e,e),$$

where $F(g,h)=\sum_{n\geq 1}f^{(n)}(g,h)$ with $f^{(n)}(g,h)$ being the probability of reaching h for the first time after n steps when starting at g. If p is symmetric, then the (left invariant) Green distance is defined by $d_G(g,h):=-\ln F(g,h)$. The drift $\ell_{p,G}$ for that distance coincides with the entropy h_p ([**BP**], Proposition 6.2, [**BHM**]) and the volume entropy is 1, so that there is equality in (1) for that distance ([**BHM**]).

We now turn to another representation of the drift and entropy. Let X be a compact space. X is called a G-space if the group G acts by homeomorphisms on X. This action extends naturally to probability measures on X. We say that the measure ν on X is stationary if $\sum_{g\in G}(g_*\nu)p(g)=\nu$. The *entropy* of a stationary

measure ν is defined by

$$h_p(X,\nu) := -\sum_{g\in G}\left(\int_X \ln \frac{dg_*^{-1}\nu}{d\nu}(\xi)d\nu(\xi)\right)p(g). \tag{3}$$

The entropy h_p and the linear drift ℓ_p are given by variational formulas over stationary measures (see [**KV**] for the entropy, [**KL**] for the linear drift):

$$h_p = \max\{h_p(X,\nu); X \text{ } G\text{-space and } \nu \text{ stationary on } X\}, \tag{4}$$

$$\ell_p = \max\Big\{\sum_{g\in G}\left(\int_{\overline{G}}\xi(g^{-1})d\nu(\xi)\right)p(g); \nu \text{ stationary on } \overline{G}\Big\}, \tag{5}$$

where $\overline{G}$ is the Busemann compactification of G, the elements of which are horofunctions ξ on G. A pair (X,ν), where X is a G-space and ν a p-stationary measure is called a *boundary* if, for P-a.e.ω, $(X_n(\omega))_*\nu$ converge towards a Dirac measure. It is called a *Poisson boundary* if it is a boundary and it realizes the maximum in formula (4).

From the definition of ℓ_p and h_p, one sees that the mappings $p\mapsto \ell_p$ and $p\mapsto h_p$ are uppersemicontinuous on $\mathcal{P}(B)$. Erschler ([**Er**]) raised the question of continuity of these functions and gave examples where these mappings are not continuous on the closure of $\mathcal{P}(B)$. The question of continuity in general on the interior of $\mathcal{P}(B)$ is open. In the rest of the paper, we discuss several examples where one can prove stronger regularity results.

2. Nearest neighbour random walks on a free group

In the case when the group G is a *free group* with d generators, $d\geq 2$, and p is supported by these generators, explicit computations can be made (see [**DM**]).

Let G be the free group with set of generators $S=\{\pm i; i=1,\dots,d\}$, where $-i=i^{-1}$ for $i\in S$. Let $\mathcal{P}(S)$ be the set of probability measures on G with support S. Since $d\geq 2$, as n goes to infinity, the reduced word representing $X_n(\omega)$ converges towards an infinite reduced word $X_\infty(\omega)=s_1(\omega)s_2(\omega)\cdots$ with $s_j(\omega)\neq -s_{j+1}(\omega)$. Denote by G_∞ the space of infinite reduced words. The stationary measure is unique: it is the distribution ν of $X_\infty(\omega)$. Then (G_∞,ν) is the Poisson boundary and G_∞ is the Busemann boundary of G. Let $q_i=P\big(\{\omega; s_1(\omega)=i\}\big)=\nu([i])$, where $[i]$ consists of all infinite words in G_∞ starting with letter $i\in S$. We have $\sum_{i\in S}q_i=1$. Let $i_1\dots i_k$ be a reduced word in G. Then ν is uniquely determined by the values $\nu([i_1\dots i_k])=F(e,i_1\dots i_k)(1-q_{-i_k})$, where $F(e,i_1\dots i_k)$ is the probability of hitting $i_1\dots i_k$ when starting at the identity e. Formula (5) writes:

$$\ell_p = 1-2\sum_{i\in S}p_iq_{-i},$$

where $p_i=p(i)$. In order to write the formula for the entropy, we introduce $z_i:=F(e,i)$ for $i\in S$. The density $\dfrac{dg_*^{-1}\nu}{d\nu}(\xi)$ gives the minimal positive harmonic function with pole at $\xi=i_1i_2\dots\in G_\infty$. The Green function satisfies the following multiplicative structure:

$$G(e,i_1\dots i_k)=F(e,i_1)G(i_1,i_1\dots i_k)=F(e,i_1)G(e,i_2\dots i_k).$$

This yields together with [**L1**, Theorem 2.10]

$$\frac{di_*^{-1}\nu}{d\nu}(\xi) \;=\; \lim_{k\to\infty}\frac{G(-i,i_1\dots i_k)}{G(e,i_1\dots i_k)} = \begin{cases} z_i, & \text{if } i_1\neq -i,\\ z_{-i}^{-1}, & \text{if } i_1=-i.\end{cases}$$

Formula (4) writes:

$$h_p = \sum_{i\in S} p_i\big[q_{-i}\ln z_{-i} - (1-q_{-i})\ln z_i\big].$$

We can express the q_i in terms of the p_i, and vice versa, thanks to the *traffic equations*: using the Markov property, we can write:

$$z_i \;=\; p_i + z_i\sum_{j\in S\setminus\{i\}} p_j z_{-j} \quad\text{and}\quad q_i \;=\; z_i(1-q_{-i}).$$

The last equation arises from the formula given for $\nu([i])$ above. Setting $Y := \sum_{j\in S} p_j z_{-j}$, we get

$$p_i = \frac{z_i(1-Y)}{1-z_iz_{-i}} \quad\text{and}\quad z_i \;=\; \frac{q_i}{1-q_{-i}},$$

so that we find:

$$\ell_p \;=\; 1-\frac{2}{A}\sum_{i\in S}\frac{q_iq_{-i}(1-q_i)}{1-q_i-q_{-i}}, \quad\text{where}\quad A \;=\; (1-Y)^{-1} \;=\; 1+\sum_{i\in S}\frac{q_iq_{-i}}{1-q_i-q_{-i}},$$

which writes:

$$\ell_p = \frac{B}{A},\ \text{where } B := 1-\sum_{i\in S}\frac{q_iq_{-i}(1-2q_i)}{1-q_i-q_{-i}}.$$

Hence, in terms of the $q_i, i\in S$, p_i and ℓ_p are rational, and the expression of h_p involves rational functions and $\ln q_i, \ln(1-q_i)$.

Proposition 2.1. *The mappings $p\mapsto \ell_p$ and $p\mapsto h_p$ are real analytic on $\mathcal{P}(S)$.*

Proof. Since all formulas are explicit in terms of the q_i, we only have to check that the q_i are real analytic functions on $\mathcal{P}(S)$. First, since z_i can be seen as a sum of probabilities of different paths in G we can write z_i as a power series in terms of the p_i's and the additional variable $z\in\mathbb{C}$ (contained in a suitable domain of convergence), namely as

$$z_i(z) = \sum_{(n_1,\dots,n_{2d})\in\mathbb{N}^{2d}} c(n_1,\dots,n_{2d})p_1^{n_1}p_{-1}^{n_2}p_2^{n_3}p_{-2}^{n_4}\cdot\ldots\cdot p_d^{n_{2d-1}}p_{-d}^{n_{2d}}z^{n_1+\dots+n_{2d}},$$

where $c(n_1,\dots,n_{2d})\geq 0$. Note that $z_i = z_i(1)$. Since the spectral radius is strictly smaller than 1 (see e.g. [**W**, Cor. 12.5]), the power series $G(e,i|z) = \sum_{n\geq 0} p^{(n)}(i)z^n$ has radius of convergence strictly bigger than 1 and satisfies $G(e,i|z)\geq z_i(z)$ for all real $z>0$. That is, for each $p\in\mathcal{P}(S)$, $z_i(z)$ has radius of convergence $R_i>1$. Choose now any $\delta>0$ with $1+\delta<R_i$. Then

$$\begin{aligned} z_i \;&=\; z_i(1)\leq z_i(1+\delta)\\ &=\sum_{(n_1,\dots,n_{2d})\in\mathbb{N}^{2d}} c(n_1,\dots,n_{2d})\big((1+\delta)p_1\big)^{n_1}\cdot\ldots\cdot\big((1+\delta)p_{-d}\big)^{n_{2d}}<\infty.\end{aligned}$$

In other words, $z_i = z_i(1)$ is real analytic in a neighbourhood of any $p \in \mathcal{P}(S)$. The equations $q_i = z_i(1-q_{-i}), q_{-i} = z_{-i}(1-q_i)$ give

$$q_i = \frac{z_i(1-z_{-i})}{1-z_i z_{-i}},$$

and this finishes the proof. □

Observe that, for $d = 1$, the group G is $\mathbb{Z}$, $S = \{\pm 1\}$ and $p \mapsto \ell_p = |p_1 - p_{-1}|$ is not a real analytic function on $\mathcal{P}(\pm 1)$.

The formulas are even simpler when the probability p is symmetric. Let $\mathcal{P}_\sigma(S)$ be the set of symmetric probability measures on S; elements of $\mathcal{P}_\sigma(S)$ are described by d positive numbers $\{p_1, \cdots, p_d\}$ such that $\sum_{i=1}^d p_i = 1/2$. If $p \in \mathcal{P}_\sigma(S)$, $q_i = q_{-i}$ and we have:

$$\begin{aligned}
\ell_p &= \frac{B}{A} \text{ with } A = 1 + 2\sum_{i=1}^d \frac{q_i^2}{1-2q_i} \text{ and } B = 1 - 2\sum_{i=1}^d q_i^2, \\
h_p &= -\frac{2}{A}\sum_{i=1}^d q_i(1-q_i)\ln\frac{q_i}{1-q_i}, \text{whereas} \\
p_i &= \frac{q_i(1-q_i)}{A(1-2q_i)} \quad \text{for } i = 1, \ldots, d.
\end{aligned}$$

Proposition 2.2. *The functions $p \mapsto \ell_p$ and $p \mapsto h_p$ reach their maxima on $\mathcal{P}_\sigma(S)$ at the constant vector $p_0 = (1/2d, \ldots, 1/2d)$ and*

$$\ell_{p_0} = 1 - \frac{1}{d}, \quad h_{p_0} = \big(1 - \frac{1}{d}\big)\ln(2d-1).$$

Proof. By symmetry, the constant vector p_0 is a critical point for ℓ_p. At p_0, $q_i = 1/2d$ by symmetry and $\ell_{p_0} = 1 - 1/d, h_{p_0} = (1-1/d)\ln(2d-1)$ by the formulas above (observe that these expressions are also valid for $d = 1$: the only point of $\mathcal{P}_\sigma(\pm 1)$ is $(1/2, 1/2)$, for which $\ell = 0 = 1 - 1/d$ and $h = 0 = (1-1/d)\ln(2d-1)$). Moreover, the volume entropy of G is $\ln(2d-1)$. By (1), the result for ℓ_p implies that $(1-1/d)\ln(2d-1)$ is the maximal value that the entropy might take on $\mathcal{P}_\sigma(S)$. Since this is the entropy h_{p_0}, p_0 achieves the maximum of the entropy as well.

We are going to prove that the function $(q_1, \ldots, q_d) \mapsto B/A$ has a unique critical point on the set $\{(q_1, \ldots, q_d); q_j > 0, \sum_{j=1}^d q_j = 1/2\}$. Observe that the formula for ℓ_p is continuous on the domain $0 \le q_i \le 1/2$ and that the value of ℓ_p at the boundary of the domain $\{(q_1, \ldots, q_d); q_j > 0, \sum_{j=1}^d q_j = 1/2\}$ is the one computed with only the non-zero q_i's on a free group with a smaller set of generators. Since at the constant vector p_0, $\ell_{p_0} = 1 - 1/d$, it follows, by induction on the dimension, that the critical point p_0 is a maximum. The proof for $d = 2$ is the same as in the general case: there is only one critical point by the argument below and the limit of the expression for ℓ_p at $(0, 1/2), (1/2, 0)$ is 0.

Using a Lagrange multiplier, we are looking for the critical points of the function $F(q, \lambda) = \ell_p - \lambda(\sum_{j=1}^d q_j - 1/2)$ satisfying $0 \le q_j \le 1/2$ for $j = 1, \ldots, d$. Setting as above

$$A = 1 + 2\sum_{i=1}^d \frac{q_i^2}{1-2q_i} \quad \text{and} \quad B = 1 - 2\sum_{i=1}^d q_i^2,$$

all equations $\frac{\partial F}{\partial q_i} = 0$ depend only on A, B, λ and q_i.

Indeed, they write $G(A, B, \lambda, q_i) = 0$, where:

$$G(A, B, \lambda, q) = 16Aq^3 + 4q^2(\lambda A^2 - 4A - B) + 4q(-\lambda A^2 + A + B) + \lambda A^2.$$

If, for fixed A, B, λ, the equation $G(A, B, \lambda, q_i) = 0$ has only one solution $q \in [0, 1/2)$, then, for these values of A, B, λ, the only possible critical point of F is $q_j = 1/2d$ for all j. Then, unless $A = \frac{2d-1}{2d-2}, B = \frac{2d-1}{2d}$, there is no critical point for F with those values of A, B.

To summarize, we only have to verify that the equation $G(A, B, \lambda, q_i) = 0$ has at most one solution $q \in [0, 1/2)$ for all A, B, λ with $0 < B < 1 < A$.

The function $q \mapsto G(A, B, \lambda, q)$ is a third degree polynomial with positive highest coefficient, $1/2$ is a critical point and $G(A, B, \lambda, 1/2) = B > 0$. Therefore, there is at most one solution $q \in [0, 1/2)$. □

Let us mention the recent article [**GMM**, Corollary 2.7], where bounds for ℓ_p and h_p are provided which are equal to the values of ℓ_{p_0} and h_{p_0} in our context of free groups. It is likely that p_0 gives also the maximum of the entropy on the whole $\mathcal{P}(S)$, but we do not have a proof of that fact. We also conjecture that the mapping $p \mapsto \ell_p$ is a concave function; numerical computer calculations of the drift for small $d \in \mathbb{N}$ supports and confirms this conjecture, but we do not have a proof for general d.

3. Free products, Artin dihedral groups and braid groups

The computations in Section 2 have been known for fifty years (even if Proposition 2.2 seems to be formally new). There are few other examples where it is possible to describe geometrically the Poisson boundary and the Busemann boundary, and it is even rarer to be able to give useful formulas for the stationary measure. In this section, we review the examples we are aware of.

One important concept of constructing new groups from given ones is the free product of groups. The crucial point is that free products have a tree-like structure. More precisely, suppose we are given finitely generated groups $G_1, \dots, G_r$ equipped with finitely supported probability measures $p_1, \dots, p_r$. The identity of G_i is denoted by e_i, and w.l.o.g. we assume that these groups are pairwise disjoint and we exclude the case $r = 2 = |G_1| = |G_2|$ (this case leads to recurrent random walks in our setting). The free product $G_1 * \cdots * G_r$ is given by

$$G = *_{i=1}^r G_i = \big\{x_1 x_2 \dots x_n \big| x_j \in \bigcup_{i=1}^r G_i \setminus \{e_i\}, x_j \in G_k \Rightarrow x_{j+1} \notin G_k\big\} \cup \{e\},$$

the set of finite words over the alphabet $\bigcup_{i=1}^r G_i \setminus \{e_i\}$ such that two consecutive letters do *not* come from the same group G_k, where e describes the empty word. A group operation on G is given by concatenation of words with possible contractions and cancellations in the middle such that one gets a reduced word as above. For $x = x_1 \dots x_n \in G$, define the *block length* of x as $\|x\| := n$. Let us mention that the free group with d generators is a free product $\mathbb{Z} * \cdots * \mathbb{Z}$ with k free factors $\mathbb{Z}$.

A random walk on G is constructed in a natural way as follows: we lift p_i to a probability measure $\bar{p}_i$ on G: if $x = x_1 \dots x_n \in G$ with $x_n \notin G_i$ and $v, w \in G_i$, then $\bar{p}_i(xv, xw) := p_i(v, w)$. Otherwise we set $\bar{p}_i(x, y) := 0$. Choose $0 < \alpha_1, \dots, \alpha_r \in \mathbb{R}$

with $\sum_{i=1}^r \alpha_i = 1$. Then we obtain a new probability measure on G defined by

$$p = \sum_{i=1}^r \alpha_i \bar{p}_i$$

with $B = \operatorname{supp}(p) = \bigcup_{i=1}^r \operatorname{supp}(p_i)$. We consider random walks $(X_n)_{n\in\mathbb{N}_0}$ on G starting at e, which are governed by p. For $i \in \{1,\dots,r\}$, denote by ξ_i the probability of hitting the set $G_i \setminus \{e_i\}$ when starting at e. The spectral radius $\varrho(p)$ is strictly less than 1 due to the non-amenability of G. Let ∂G_i be the Martin boundary of G_i with respect to p_i, and denote by G_∞ the set of infinite words $x_1x_2\dots$ such that $x_i \in G_k$ implies $x_{i+1} \notin G_k$. Then the Martin boundary of G is given by

$$\partial G = G_\infty \cup \bigcup_{i=1}^r \{x\xi; x = x_1\dots x_n \in G, x_n \notin G_i, \xi \in \partial G_i\};$$

see e.g. [**W**, Proposition 26.21]. The random walk on G converges almost surely to an infinite word in G_∞ in the sense that the common prefix of the word at time n and the infinite limit word tends to infinity as $n \to \infty$. The limit distribution ν is determined by

$$\nu\big(\{x_1x_2\dots \in G_\infty; x_1 = y_1, \dots, x_n = y_n\}\big) = F(e, y_1\dots y_n)\big(1 - (1-\xi_i)G_i(\xi_i)\big),$$

where $n \in \mathbb{N}$, $y_1\dots y_n \in G$ with $y_n \notin G_i$, $F(e, y_1\dots y_n)$ being the probability of hitting $y_1\dots y_n$ and $G_i(z) = \sum_{n\geq 0} p_i^{(n)}(e_i)z^n$ with $z \in \mathbb{C}$; see e.g. [**Gi1**].

The next propositions summarize results about regularity of drift and entropy. Explicit formulas can be found in the cited sources.

Proposition 3.1 ([**Gi1**])**.** *The drift w.r.t. the block length $\ell_B = \lim_{n\to\infty} \frac{1}{n}\|X_n\|$ exists and varies real analytically in $p \in \mathcal{P}(B)$.*

Proof. In [**Gi1**, Equ. (9)] a formula for ℓ_B is given:

$$\ell_B = \sum_{i=1}^r \alpha_i \frac{1-\xi_i}{\xi_i}\big(1 - (1-\xi_i)\,G_i(\xi_i)\big).$$

Let be $d = |B| - 1$, and write $p = (q_1, \dots, q_d) \in \mathcal{P}(B)$. Analogously to the proof of Proposition 2.1 one can write ξ_i as a power series (evaluated at $z = 1$) in the form

$$\xi_i(z) = \sum_{(n_1,\dots,n_d)\in\mathbb{N}^d} c(n_1,\dots,n_d) q_1^{n_1} q_2^{n_2} \dots q_d^{n_d} z^{n_1+\dots+n_d}, \quad z \in \mathbb{C}.$$

Since $\varrho(p) < 1$ the Green functions $G(g|z) = \sum_{n\geq 0} p^{(n)}(g)z^n$, $g \in G$, have radii of convergence $R = 1/\varrho(p) > 1$ and dominate $\xi_i(z)$ for real $z > 0$. Hence, $\xi_i(z)$ has radius of convergence bigger than 1, which in turn – following the same argumentation as in Proposition 2.1 – yields real analyticity of $\xi_i = \xi_i(1)$ in a neighbourhood of any $p \in \mathcal{P}(B)$. Furthermore, $G_i(z)$ can be expanded in the same form as $\xi_i(z)$ and, for each real positive $z_0 < 1$, the mapping $p \mapsto G_i(z_0)$ is real analytic. Since $\xi_i < 1$ (see e.g. [**Gi1**, Lemma 2.3]) the mapping $p \mapsto G_i(\xi_i)$ is also real analytic as a composition of real analytic functions. This yields the proposed statement. □

Proposition 3.2 ([**Gi1**])**.** *Let p govern a nearest neighbour random walk on G, that is, the length $|g|$ is computed with respect to the generator B. Then the drift function $p \mapsto \ell_p = \lim_{n\to\infty} \frac{1}{n}|X_n|$ is real analytic.*

Proof. By the formula for ℓ given in [**Gi1**, Section 7] we just have to check that the mapping

$$p \mapsto \widetilde{G}_j(y,z) := \sum_{m,n\geq 0} \sum_{x\in G_j:|x|=m} p_j^{(n)}(e_j,x) y^n z^m$$

is real analytic for all $y \in (0,1)$ and $z = 1$. For a moment fix $y < 1$ and choose $\delta > 0$ small enough such that $y(1+2\delta)^2 < 1$. Since $p_j^{(n)}(e_j,x) > 0$, $x \in G_j$ with $|x| = m$, implies $n \geq m$, we get

$$\widetilde{G}_j\big(y,(1+2\delta)^2\big) = \sum_{m,n\geq 0} \sum_{x\in G_j:|x|=m} p_j^{(n)}(e_j,x)\big(y(1+\delta)^2\big)^n \leq \frac{1}{1-y(1+2\delta)^2} < \infty.$$

This yields $\frac{\partial}{\partial z}\widetilde{G}\big(y,(1+\delta)^2\big) < \infty$. Since each term $p_j^{(n)}(e_j,x)$ can be written as a polynomial

$$\sum_{\substack{(n_1,\dots,n_d)\in\mathbb{N}^d:\\ n_1+\dots+n_d=n}} c(n_1,\dots,n_d) q_1^{n_1}\cdot\ldots\cdot q_d^{n_d},$$

where $p = (q_1,\dots,q_d) \in \mathcal{P}(B)$ and $c(n_1,\dots,n_d) \geq 0$, we rewrite $\frac{\partial}{\partial z}\widetilde{G}\big(y,(1+\delta)^2\big)$ as

$$\frac{1}{1+\delta} \sum_{m,n\geq 0} \sum_{x\in G_j:|x|=m} m\big(p_j^{(n)}(e_j,x)(1+\delta)^n\big)\big(\xi_j(1+\delta)\big)^n.$$

That is, $\sum_{m,n\geq 0}\sum_{x\in G_j:|x|=m} m p_j^{(n)}(e_j,x)\xi_j^n$ is real analytic in $\mathcal{P}(B)$ as a composition of real analytic functions, and this yields the claim. □

Let us mention that – in contrast to Proposition 2.2 – simple random walk (that is, μ is the equidistribution on $\mathrm{supp}(\mu)$) is not necessarily the fastest random walk. Namely, it can be verified with the help of MATHEMATICA that – with p_i describing the simple random walk on G_i – the simple random walk on $(\mathbb{Z}/3\mathbb{Z}) * (\mathbb{Z}/2\mathbb{Z})$ (that is, $\alpha_1 = 2/3, \alpha_2 = 1/3$) is slower than the random walk on this free product with the parameters $\alpha_1 = \alpha_2 = 1/2$.

Furthermore, we have the following regularity result:

Proposition 3.3 ([**Gi3**])**.** *Assume that $h_i := -\sum_{g\in G_i} p_i(g)\ln p_i(g) < \infty$ for all $i \in \{1,\dots,r\}$, that is, all random walks on the factors G_i have finite single-step entropy. Then the mapping $p \mapsto h_p$ is real analytic.*

We now turn to another class of groups whose Cayley graphs have a tree-like structure. A group G is called *virtually free* if it has a free subgroup of finite index. At this point we assume that G has a free subgroup with at least $d \geq 2$ generators; otherwise, G is a finite extension of $\mathbb{Z}$ where we either get recurrent random walks or non-regularity points on $\mathbb{P}(B)$. It is well-known that virtually free groups can be constructed from a finite number of finite groups by iterated amalgamation and HNN extensions. Each element of G can be written as $x_1\dots x_n h$, where $x_i \in \{\pm i; i = 1,\dots d\}$ and h being one of finitely many representatives for the different cosets. Suppose we are given a weight or length function $l(\pm i) \in \mathbb{R}$ for $i \in \{1,\dots,d\}$. Then a natural length function on G is defined by $l(x_1\dots x_n h) = \sum_{j=1}^n l(x_j)$. We have the following result:

Proposition 3.4 ([**Gi2**])**.** *Let G be a virtually free group. Let p govern a finite range random walk on G. Then the mapping $p \mapsto \lim_{n\to\infty} l(X_n)/n$ is real analytic.*

Proof. Random walks on virtually free groups can be interpreted as a random walk on a regular language in the sense of [**Gi2**]. The claim follows from the formula for $\lim_{n\to\infty} l(X_n)/n$, the drift with respect to the length function l, given in [**Gi2**, Theorem 2.4]. Due to non-amenability of G we have again $\varrho(p) < 1$. The rest follows analogously as in the proofs of Propositions 2.1 and 3.1. □

For the special case l being the natural word length the last proposition is also covered by Corollary 4.2.

At this point we want to mention the article [**MM2**], which uses similar techniques to establish statements about the drift of random walks on the braid group B_3 and on Artin groups of dihedral type. Traffic equations are established, whose unique solutions lead to formulas for the drift. For random walks on these groups there might occur transitions (when varying the probability measures of constant support), where one has no regularity. An explicit example for non-differentiability points is given on the braid group B_3; at these points the random walk changes its behaviour essentially in terms of how the limit words looks like. However, [**MM2**] gives explicit formulas for the drift in terms of the solutions of the traffic equations splitted up into different branches. By methods similar to the above, one can show that the drift is real analytic on each branch. Indeed, solutions of the traffic equations can be written as converging power series as in the proofs of Propositions 3.1 and 3.2.

4. Hyperbolic groups

A geodesic metric space is called *hyperbolic* if geodesic triangles are thin: there is $\delta \geq 0$ such that each side of a geodesic triangle is contained in a δ-neighbourhood of the union of the other two sides. A finitely generated group is called hyperbolic if the Cayley graph defined by some finite symmetric generating set is hyperbolic. This property does not depend on the choice of a generating set. Free groups are hyperbolic, as are fundamental groups of compact manifolds of negative curvature, and small cancellation groups. See e.g. [**GH**] for the main geometric properties of hyperbolic groups. The geometric boundary of a hyperbolic space is the space of equivalence classes of geodesic rays, where two geodesic rays are equivalent if they are at a bounded Hausdorff distance. The geometric boundary ∂G of the Cayley graph of a hyperbolic group G is a compact G-space. It is endowed with the Gromov metric (see [**GH**]). The mapping $\Phi : G \to \mathbb{Z}^G, \Phi(g)(h) = |h^{-1}g| - |g|$ is an injective mapping such that $\Phi(G)$ is relatively compact for the product topology. The Busemann compactification $\overline{G}$ is the closure of $\Phi(G)$ in $\mathbb{Z}^G$. There is an equivariant homeomorphism $\pi : \overline{G} \setminus G \to \partial G$ (see e.g. [**WW**]). The homeomorphism π is finite-to-one (see e.g. [**CP**]). Following [**Bj**], we say that G with the generating set S satisfies the *basic assumption* (BA) if the homeomorphism π is one-to-one. In this case, we write, for $\xi \in \partial G, h \in G$, $\xi(h)$ for the value at h of the sequence $\pi^{-1}\xi \in \mathbb{Z}^G$. Free groups and surface groups with their natural generators satisfy (BA). It is an open problem whether any hyperbolic group admits a symmetric generating set with the property (BA).

Let p be a probability measure on G with finite support. Then, there is a unique p-stationary probability measure ν_p on ∂G and $(\partial G, \nu_p)$ is a Poisson boundary for (G, p)([**An**], [**K**] Theorem 7.6). If (BA) is satisfied, the measure ν_p is the unique stationary probability measure on the Busemann compactification and formulas (4)

and (5) write:
(6)

$$h_p \;=\; -\sum_{g\in B}\left(\int_{\partial G}\ln\frac{dg_*^{-1}\nu_p}{d\nu_p}(\xi)d\nu_p(\xi)\right)p(g),\quad \ell_p \;=\; \sum_{g\in B}\left(\int_{\partial G}\xi(g^{-1})d\nu_p(\xi)\right)p(g).$$

Proposition 4.1. *Assume that (G,S) is a non-elementary hyperbolic group and satisfies (BA). Let $p\in\mathcal{P}(B)$, α be small enough, and let f be an α-Hölder continuous function on ∂G. Then the mapping $p\mapsto\int_{\partial G}f(\xi)d\nu_p(\xi)$ is real analytic on a neighbourhood of p in $\mathcal{P}(B)$.*

Proof. Let $\mathcal{K}_\alpha$ be the space of α-Hölder continuous functions on ∂G. The space $\mathcal{K}_\alpha$ is a Banach space with norm $\|f\|_\alpha$, where

$$\|f\|_\alpha \;=\; \max_{\xi\in\partial G}|f(\xi)|+\sup_{\xi,\eta\in\partial G:\xi\neq\eta}\frac{|f(\xi)-f(\eta)|}{(d(\xi,\eta))^\alpha}.$$

For $p\in\mathcal{P}(B)$, let $\mathcal{Q}_p$ be the operator on $\mathcal{K}_\alpha$ defined by

$$\mathcal{Q}_pf(\xi) \;=\; \sum_{g\in B}f(g^{-1}\xi)p(g).$$

Clearly, the mapping $p\mapsto\mathcal{Q}_p$ is real analytic from $\mathcal{P}(B)$ into $\mathcal{L}(\mathcal{K}_\alpha)$. If G is not elementary and satisfies (BA), it can be shown (see [**Bj**], Lemma 4) that, for α small enough, $f\mapsto\int fd\nu_p$ is an isolated eigenvector for the transposed operator $\mathcal{Q}_p^*$ on the dual space $\mathcal{K}_\alpha^*$. The proposition follows by a perturbation lemma. □

Corollary 4.2. *Assume that (G,S) is a non-elementary hyperbolic group and satisfies (BA). Then the mapping $p\mapsto\ell_p$ is real analytic.*

Indeed, the function $\xi(g^{-1})$ in formula (6) belongs to $\mathcal{K}_\alpha$ for all α. Corollary 4.2 is due to [**L2**] in the case of the free group. P. Mathieu ([**Ma2**]) proved the C^1 regularity and gave a formula for $\nabla_p\nu_p$ and $\nabla_p\ell_p$ in the symmetric case, to be compared with formulas for linear response of dynamical systems (cf. [**R**]).

The formula (6) for the entropy is valid in general, even without the (BA) hypothesis, but observe that the integrand $\varphi_p(g,\xi):=-\ln\frac{dg_*^{-1}\nu_p}{d\nu_p}(\xi)$ is itself a function of p. To study this function, we use the description by A. Ancona ([**An**]) of the Martin boundary of a random walk with finite support on a hyperbolic group. Recall that $F_p(g,h)$ is the probability of reaching h starting from g in dependence of p.

Proposition 4.3 ([**An**])**.** *Assume that G is hyperbolic and that p has finite support. Then*

$$\varphi_p(g,\xi) \;=\; \lim_{h\to\xi}\ln\frac{F_p(e,h)}{F_p(g^{-1},h)}\qquad \textit{for all } g\in G,\xi\in\partial G.$$

A consequence of the proof of Proposition 4.3 is that, for all $g\in G$, for α small enough $\varphi_p(g,\xi)\in\mathcal{K}_\alpha$ (see [**INO**]). In the case of free groups, Proposition 4.3 goes back to Derriennic ([**De1**]) and using his arguments one can prove:

Proposition 4.4 ([**L2**])**.** *If G is a free group and p has finite support B, there is α small enough that, for all $g\in B$, the mapping $p\mapsto\varphi_p$ is real analytic from a neighbourhood of p in $\mathcal{P}(B)$ into $\mathcal{K}_\alpha$.*

Corollary 4.5 ([**L2**]). *If G is a free group and p has finite support B, the mapping $p \mapsto h_p$ is real analytic on $\mathcal{P}(B)$.*

For cocompact Fuchsian groups there is the following recent result, to be compared with Corollary 4.2:

Proposition 4.6 ([**HMM**]). *Let G be a cocompact Fuchsian group with planar presentation. Then the mapping $p \mapsto \ell_p$ is real analytic.*

For a general hyperbolic group, we have a weaker result:

Proposition 4.7 ([**L3**]). *If G is a hyperbolic group and p has finite support B, there is α small enough such that, for all $g \in B$, the mapping $p \mapsto \varphi_p$ is Lipschitz continuous from a neighbourhood of p in $\mathcal{P}(B)$ into $\mathcal{K}_\alpha$.*

Corollary 4.8 ([**L3**]). *If G is a hyperbolic group and p has finite support B, the mappings $p \mapsto h_p$, $p \mapsto \ell_p$ are Lipschitz continuous on $\mathcal{P}(B)$.*

[**Gi4**] proves also regularity properties of the mapping $p \mapsto h_p$ for random walks on regular languages, which adapt, for instance, to the case of virtually free groups.

The best results of regularity to-date are due to P. Mathieu:

Proposition 4.9 ([**Ma2**]). *If G is a hyperbolic group and B is finite and symmetric, the mapping $p \mapsto h_p$ is C^1 on the set $\mathcal{P}_\sigma(B)$ of symmetric probability measures on B, endowed with its natural smooth structure.*

Let $\lambda \mapsto p_\lambda, \lambda \in [-\varepsilon, +\varepsilon]$ be a smooth curve in $\mathcal{P}(B)$. We write:

$$\begin{aligned}
& \lim_{\lambda\to 0} \frac{h(p_\lambda) - h(p_0)}{\lambda} \\
= \; & \lim_{\lambda\to 0} \frac{1}{\lambda} \sum_{g\in B} \left(\int_{\partial G} \varphi_{p_\lambda}(g,\xi) d\nu_{p_\lambda}(\xi) p_\lambda(g) - \int_{\partial G} \varphi_{p_0}(g,\xi) d\nu_{p_0}(\xi) p_0(g) \right) \\
= \; & \lim_{\lambda\to 0} \frac{1}{\lambda} \sum_{g\in B} \left(\int_{\partial G} (\varphi_{p_\lambda}(g,\xi) - \varphi_{p_0}(g,\xi)) d\nu_{p_\lambda}(\xi) \right) p_\lambda(g) + \\
& \quad + \sum_{g\in B} \left(\int_{\partial G} \varphi_{p_0}(g,\xi)) [\lim_{\lambda\to 0} \frac{1}{\lambda} (d\nu_{p_\lambda} - d\nu_{p_0})](\xi) \right) p_\lambda(g) + \\
& \quad\quad + \sum_{g\in B} \left(\int_{\partial G} \varphi_{p_0}(g,\xi)) d\nu_{p_0}(\xi) \right) [\lim_{\lambda\to 0} \frac{1}{\lambda} (p_\lambda(g) - p_0(g)].
\end{aligned}$$

The third line converges by definition. To prove that the second line converges, P. Mathieu observes that the Green metric $-\ln F_{p_0}(g,h)$ on G satisfies (BA) and a form of hyperbolicity that allows him to extend Proposition 4.1. More precisely, he shows directly the differentiability of $\lambda \mapsto \int f(\xi) d\nu_{p_\lambda}(\xi)$, for $f \in \mathcal{K}_\alpha$, and gives a formula for the derivative. For the first line, P. Mathieu shows a general result for any non-amenable group G and p_λ with finite support. In our case, his result writes:

Proposition 4.10 ([**Ma2**]). *Let $\lambda \mapsto p_\lambda, \lambda \in [-\varepsilon, +\varepsilon]$ be a smooth curve in $\mathcal{P}(B)$. Then,*

$$\lim_{\lambda\to 0} \frac{1}{\lambda} \sum_{g\in B} \left(\int (\varphi_{p_\lambda}(g,\xi) - \varphi_{p_0}(g,\xi)) d\nu_{p_\lambda}(\xi) \right) p_\lambda(g) \;=\; 0.$$

It is likely that the function $p \mapsto h_p$ has more regularity on $\mathcal{P}(B)$, but this is an open problem.

Another natural extension is towards more general families of probability measures on G. Proposition 2.1 is valid for p varying in finite dimensional affine subsets of $\{p; \sum_{g\in G} e^{\gamma|g|}p(g) < +\infty\}$ for some $\gamma > 0$ (see [**L1**]). The other properties rest on Harnack inequality at infinity (see [**An**]), which has been proven only for probability measures with finite support on hyperbolic groups. Finally, let $\mathcal{P}^1(G)$ be the set of probabilities on G satisfying $\sum_{g\in G} |g|p(g) < +\infty$ endowed with the topology of convergence on the functions which grow slower than $C|g|$ at infinity. The first observation on this topic of regularity of the entropy is the fact that, if G is hyperbolic, $p \mapsto h_p$ is continuous on $\mathcal{P}^1(G)$ ([**EKc**]).

References

[**An**] A. Ancona, Théorie du potentiel sur les graphes et les variétés, École d'été de Saint-Flour XVIII,1988, *Lecture Notes in maths,* **1427** (1990) 1–112.

[**Av**] A. Avez, Entropie des groupes de type fini, *C. R. Acad. Sc. Paris Sér A-B* **275** (1972), A1363–A1366.

[**Bj**] M. Bjorklund, Central Limit Theorem for Gromov Hyperbolic Groups, *J. Theo. Probability*, **23** (2010), 871–887.

[**BHM**] S. Blachère, P. Haïssinsky and P. Mathieu, Asymptotic entropy and Green speed for random walks on countable groups, *Ann. Prob.,* **36** (2008), 1134–1152.

[**BP**] I. Benjamini and Y. Peres, Tree-indexed random walks on groups and first passage percolation, *Probab. Theory Relat. Fields,* **98** (1994), 91–112.

[**CP**] M. Coornaert and A. Papadopoulos, Horofunctions and symbolic dynamics on Gromov hyperbolic groups, *Glasgow Math. Journal*, **43** (2001), 425–456.

[**De1**] Y. Derriennic, Marche aléatoire sur le groupe libre et frontière de Martin, *Z. Wahrscheinlichkeitstheorie verw. Geb.* **32** (1975), 261–276.

[**De2**] Y. Derriennic, Quelques applications du théorème ergodique sousadditif, *Astérisque* **74** (1980), 183–201.

[**DM**] E. B. Dynkin and M. B. Malyutov, Random walks on groups with a finite number of generators, *Dokl. Akad. Nauk SSSR*, **137** (1961), 1042–1045.

[**Er**] A. Erschler, On continuity of range, entropy and drift for random walks on groups, *in* Random Walks, Boundaries and Spectra, D. Lenz, F. Sobieszky and W. Woess ed., *Progress in Probability* bf 64, Birkhäuser Basel (2011), 55-64.

[**EKc**] A. Erschler and V. A. Kaimanovich, Continuity of entropy for random walks on hyperbolic groups, *in preparation.*

[**EKn**] A. Erschler and A. Karlsson, Homomorphisms to $\mathbb{R}$ constructed from random walks, Ann. Inst. Fourier, **60** (2010), 2095–2113.

[**GH**] É. Ghys and P. de la Harpe, Sur les groupes hyperboliques d'après Mikhael Gromov, *Progress in Mathematics* Vol.83, Birkhäuser Boston Inc., Boston, MA, 1990.

[**Gi1**] L. A. Gilch, Rate of Escape of Random Walks on Free Products, *J. of Austral. Math. Soc.,* **83** (2007) 31 - 54.

[**Gi2**] L. A. Gilch, Rate of Escape of Random walks on Regular Languages and Free Products by Amalgamation of Finite Groups, *Fifth Colloquium on Mathematics and Computer Science,* Discrete Mathematics and Computer Science Proc. AI, Assoc. Discrete Math. Theor. Comput. Sci., Nancy (2008), 405–420.

[**Gi3**] L. A. Gilch, Asymptotic entropy of random walks on free products, *Elec. J. Prob.,* **16** (2011) 76–105.

[**Gi4**] L. A. Gilch, Asymptotic entropy of random walks on regular languages, preprint, 2013.

[**GMM**] S. Gouëzel, F. Mathéus and F. Maucourant, Sharp lower bounds for the asymptotic entropy of symmetric random walks, arXiv:1209.3378, 2012.

[**Gu**] Y. Guivarc'h, Sur la loi des grands nombres et le rayon spectral d'une marche aléatoire, *Astérisque*, **74** (1980), 47–98.

[**HMM**] P. Haïssinsky, P. Mathieu and S. Müller, Renewal theory for random walks on cocompact Fuchsian groups, preprint, (2012).

[**INO**] M.Izumi, S.Neshveyev and R. Okayasu, The ratio set of the hyperbolic measure of a random walk on a hyperbolic group, *Israel J. Math.*, **163** (2008), 285–316.

[**K**] V.A. Kaimanovich, The Poisson formula for groups with hyperbolic properties, *Ann. Math.,* **152** (2000), 659–692.

[**KV**] V.A. Kaimanovich and A. M. Verschik, Random walks on discrete groups: boundary and entropy, *Annals Prob.* **11** (1983), 457–490.

[**KL**] A. Karlsson and F. Ledrappier, Drift and entropy for random walks, *Pure Appl. Math. Quarterly,* **3** (2007), 1027–1036.

[**Ki**] J.F.C. Kingman, The Ergodic Theory of Subadditive Processes, *J. Royal Stat. Soc., Ser. B,* **30** (1968), 499–510.

[**L1**] F. Ledrappier, Some Asymptotic properties of random walks on free groups, *in* Topics in Probability and Lie Groups, J.C. Taylor, ed. *CRM Proceedings and Lecture Notes,* **28** (2001), 117–152.

[**L2**] F. Ledrappier, Analyticity of the entropy for some random walks, *Groups, Geometry and Dynamics*, **6** (2012), 317–333.

[**L3**] F. Ledrappier, Regularity of the entropy for random walks on hyperbolic groups, *Annals Prob.*, to appear.

[**MM1**] J. Mairesse and F. Mathéus, Random walks on free products of cyclic groups, *J. London Math. Soc.,* **75** (2007), 47–66.

[**MM2**] J. Mairesse and F. Mathéus, Randomly growing braid on three strands and the manta ray, *Ann. Applied Proba.,* **17** (2007), 502–536.

[**Ma1**] P. Mathieu, Carne-Varopoulos bounds for centered random walks, *Annals Prob.*, **34** (2006), 987–1011.

[**Ma2**] P. Mathieu, Differentiating the entropy of random walks on hyperbolic groups, *preprint.*

[**R**] D. Ruelle, A review of linear response fo rgeneral differentiable dynamical systems, *Nonlinearity,* **22** (2009), 855–870.

[**Va**] N. Th. Varopoulos, Long range estimates for Markov chains, *Bull. Sci. Math.*, **100** (1985), 225–252.

[**V**] A. M. Verschik, Dynamic Theory of growth in groups: entropy, boundaries examples. *Russ. Math. Surveys* **55** (2000), 667–733.

[**WW**] C. Webster and A. Winchester, Boundaries of hyperbolic metric spaces, *Pacific J. Math.,* **221**(2005), 147–158.

[**W**] W. Woess, Random walks on infinite graphs and groups, Cambridge Tracts in Mathematics, vol. 138, Cambridge University Press, Cambridge, 2000.

E-mail address: gilch@TUGraz.at

E-mail address: ledrappier.1@nd.edu

Institut für Mathematische Strukturtheorie, Graz University of Technology, Steyrergasse 30/III, 8010 Graz, Austria

Department of Mathematics, University of Notre Dame, Notre Dame, IN 46556, USA